Crossword

Across

- 5. Philosopher who made scientific discoveries through careful observation and logical reasoning ... but not using scientific evidence.
- 6. Type of collision in which the colliding objects show significant deformation and generate heat. Collisions are not perfect and energy / momentum is transformed to heat or deformation of objects.
- 7. Newton's third law states that all forces react in equal and opposite.
- 8. Newton's law of motion showing that action forces on an object are balanced by reaction forces on the object that are equal and opposite.
- 9. In a closed system with perfectly elastic collisions, the momentum before a collision equals the momentum after the collision. This is of momentum.
- 10. The amount of inertia an object possesses, and depends on the amount of matter the object contains.
- 12. Inertia in motion or the product of the mass of an object times its velocity (mv). A vector quantity involving velocity, angle (direction), and mass.
- 13. Scientist who proposed 3 laws of motion: inertia, f = ma. action/reaction.

Down

- 1. Newton's law of motion showing how an object's motion does not change unless a net force acts upon it. e.g. Objects at rest stay at rest, objects in motion stay in motion unless acted upon by a net force.
- 2. Type of collision in which the colliding objects show minimal to no deformation and there is no generation of heat. Energy / momentum is totally transferred between objects.
- 3. The force acting on an object related to its mass and the acceleration due to gravity. W = mg.
- 4. Scientist who experimented (collected and tested data) that gave evidence for phenomenon in nature. Objects fall at the same rate due to gravity (without air resistance).
- 5. The "a" in f = ma. Force is related to an objects mass and "change in motion", relating to Newton's second law of motion.
- 6. Tendency of an object to resist a change in motion. Related to mass. e.g. seatbelts and airbags were developed with this in view.
- 11. Newton's law of motion for unbalanced forces acting on an object. f = ma. The acceleration of an object is equal to the net force acting on it divided by the object's mass (a = f/m).